



DMD gene

dystrophin

Normal Function

DMD, the largest known human gene, provides instructions for making a protein called dystrophin. This protein is located primarily in muscles used for movement (skeletal muscles) and in heart (cardiac) muscle. Small amounts of dystrophin are present in nerve cells in the brain.

In skeletal and cardiac muscles, dystrophin is part of a group of proteins (a protein complex) that work together to strengthen muscle fibers and protect them from injury as muscles contract and relax. The dystrophin complex acts as an anchor, connecting each muscle cell's structural framework (cytoskeleton) with the lattice of proteins and other molecules outside the cell (extracellular matrix). The dystrophin complex may also play a role in cell signaling by interacting with proteins that send and receive chemical signals.

Little is known about the function of dystrophin in nerve cells. Research suggests that the protein is important for the normal structure and function of synapses, which are specialized connections between nerve cells where cell-to-cell communication occurs.

Health Conditions Related to Genetic Changes

Duchenne and Becker muscular dystrophy

More than 2,000 mutations in the *DMD* gene have been identified in people with the Duchenne and Becker forms of muscular dystrophy. These conditions occur almost exclusively in males and are characterized by progressive muscle weakness and wasting (atrophy) and a heart condition called dilated cardiomyopathy. Most of the mutations that cause these conditions delete part of the *DMD* gene. Other mutations abnormally duplicate part of the gene or change a small number of DNA building blocks (nucleotides) in the gene.

Mutations that cause Becker muscular dystrophy, which typically has milder features and appears at a later age than Duchenne muscular dystrophy, usually lead to an abnormal version of dystrophin that retains some function. Mutations that cause the more severe Duchenne muscular dystrophy typically prevent any functional dystrophin from being produced.

Skeletal and cardiac muscle cells without enough functional dystrophin become damaged as the muscles repeatedly contract and relax with use. The damaged cells

weaken and die over time, causing the characteristic muscle weakness and heart problems seen in Duchenne and Becker muscular dystrophy.

familial dilated cardiomyopathy

X-linked dilated cardiomyopathy

More than 30 mutations in the *DMD* gene can cause an X-linked form of familial dilated cardiomyopathy. This heart condition enlarges and weakens the cardiac muscle, preventing the heart from pumping blood efficiently. Although dilated cardiomyopathy is a sign of Duchenne and Becker muscular dystrophy (described above), X-linked dilated cardiomyopathy is typically not associated with weakness and wasting of skeletal muscles.


The mutations that cause X-linked dilated cardiomyopathy preferentially affect the activity of dystrophin in cardiac muscle cells. As a result of these mutations, affected individuals typically have little or no functional dystrophin in the heart. Without enough of this protein, cardiac muscle cells become damaged as the heart muscle repeatedly contracts and relaxes. The damaged muscle cells weaken and die over time, leading to the heart problems characteristic of X-linked dilated cardiomyopathy.

The mutations that cause X-linked dilated cardiomyopathy often lead to reduced amounts of dystrophin in skeletal muscle cells. However, enough of this protein is present to prevent weakness and wasting of the skeletal muscles.

Chromosomal Location

Cytogenetic Location: Xp21.2-p21.1, which is the short (p) arm of the X chromosome between positions 21.2 and 21.1

Molecular Location: base pairs 31,119,219 to 33,339,609 on the X chromosome (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- BMD
- DMD_HUMAN
- dystrophin (muscular dystrophy, Duchenne and Becker types)

Additional Information & Resources

Educational Resources

- Madame Curie Bioscience Database: Structure and Function of the Dystrophin-Glycoprotein Complex
<https://www.ncbi.nlm.nih.gov/books/NBK6193/>

GeneReviews

- Dilated Cardiomyopathy Overview
<https://www.ncbi.nlm.nih.gov/books/NBK1309>
- Dystrophinopathies
<https://www.ncbi.nlm.nih.gov/books/NBK1119>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28DMD%5BTI%5D%29+OR+%28dystrophin%5BTI%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+720+days%22%5Bdp%5D>

OMIM

- DYSTROPHIN
<http://omim.org/entry/300377>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_DMD.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=DMD%5Bgene%5D>
- HGNC Gene Family: X-linked mental retardation
<http://www.genenames.org/cgi-bin/genefamilies/set/103>
- HGNC Gene Family: Zinc fingers ZZ-type
<http://www.genenames.org/cgi-bin/genefamilies/set/91>

- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=2928
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/1756>
- UniProt
<http://www.uniprot.org/uniprot/P11532>

Sources for This Summary

- Aartsma-Rus A, Van Deutekom JC, Fokkema IF, Van Ommen GJ, Den Dunnen JT. Entries in the Leiden Duchenne muscular dystrophy mutation database: an overview of mutation types and paradoxical cases that confirm the reading-frame rule. *Muscle Nerve*. 2006 Aug;34(2):135-44. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16770791>
- Anderson JL, Head SI, Rae C, Morley JW. Brain function in Duchenne muscular dystrophy. *Brain*. 2002 Jan;125(Pt 1):4-13. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11834588>
- Biggar WD, Klamut HJ, Demacio PC, Stevens DJ, Ray PN. Duchenne muscular dystrophy: current knowledge, treatment, and future prospects. *Clin Orthop Relat Res*. 2002 Aug;(401):88-106. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12151886>
- Cohen N, Muntoni F. Multiple pathogenetic mechanisms in X linked dilated cardiomyopathy. *Heart*. 2004 Aug;90(8):835-41. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15253946>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1768363/>
- Ehmsen J, Poon E, Davies K. The dystrophin-associated protein complex. *J Cell Sci*. 2002 Jul 15; 115(Pt 14):2801-3. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12082140>
- Ervasti JM. Dystrophin, its interactions with other proteins, and implications for muscular dystrophy. *Biochim Biophys Acta*. 2007 Feb;1772(2):108-17. Epub 2006 Jun 7. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16829057>
- Ferlini A, Sewry C, Melis MA, Mateddu A, Muntoni F. X-linked dilated cardiomyopathy and the dystrophin gene. *Neuromuscul Disord*. 1999 Jul;9(5):339-46. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/10407857>
- GeneReview: Dystrophinopathies
<https://www.ncbi.nlm.nih.gov/books/NBK1119>
- Le Rumeur E, Winder SJ, Hubert JF. Dystrophin: more than just the sum of its parts. *Biochim Biophys Acta*. 2010 Sep;1804(9):1713-22. doi: 10.1016/j.bbapap.2010.05.001. Epub 2010 May 21. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/20472103>
- Muntoni F, Torelli S, Ferlini A. Dystrophin and mutations: one gene, several proteins, multiple phenotypes. *Lancet Neurol*. 2003 Dec;2(12):731-40. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/14636778>

- Nakamura A. X-Linked Dilated Cardiomyopathy: A Cardiospecific Phenotype of Dystrophinopathy. *Pharmaceuticals (Basel)*. 2015 Jun 9;8(2):303-20. doi: 10.3390/ph8020303. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/26066469>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4491663/>
 - Neri M, Valli E, Alfano G, Bovolenta M, Spitali P, Rapezzi C, Muntoni F, Banfi S, Perini G, Gualandi F, Ferlini A. The absence of dystrophin brain isoform expression in healthy human heart ventricles explains the pathogenesis of 5' X-linked dilated cardiomyopathy. *BMC Med Genet*. 2012 Mar 28;13:20. doi: 10.1186/1471-2350-13-20.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22455600>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3331845/>
-

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/DMD>

Reviewed: February 2017
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services